

Surgical Perioperative Antibiotic Prophylaxis Recommendations

Antibiotic Choice and Duration of Perioperative Antibiotic Therapy

- **EXCLUSION CRITERIA:** Dirty or infected wounds, including presence of obvious mucopurulence, or chronically immunosuppressed patients.
- In general, antibiotic prophylaxis for clean and clean contaminated head and neck surgeries **is not indicated**.
- *In cases of multiple procedures, the procedure with the broadest antibiotic or longest duration may be appropriate.*

Recommended Antibiotic Doses

Medication	Dose	Maximum Dose	Recommended Formulation(s)
Parenteral			
Cefazolin	30 mg/kg/dose IV q8h	2000 mg (3000 mg if ≥120 kg)	IV
Clindamycin	10 mg/kg/dose IV q8h	900 mg	
Metronidazole	15 mg/kg/dose (pre-op) 10 mg/kg/dose IV q8h (post-op)	500 mg	
Enteral			
Note that doses below are given in <u>mg/kg/DAY</u> to align with <u>outpatient</u> ordering mode. Inpatient orders use <u>mg/kg/dose</u> .			
Amoxicillin	45 mg/kg/DAY divided TID	500 mg	Suspension: 80 mg/mL Tablet: 250 mg, 500 mg
Amoxicillin-clavulanate (7:1 ratio?)	45 mg/kg/DAY divided BID	875 mg	Suspension: Amoxicillin-clavulanate 400-57 mg/5mL (max at 880 mg amoxicillin/dose (11 mL) for ease of outpatient administration Tablet: Amoxicillin-clavulanate 875-125 mg
Cephalexin	75 mg/kg/DAY divided TID	500 mg	Suspension: 50 mg/mL Capsule: 250 mg, 500 mg
Clindamycin	30 mg/kg/DAY divided TID	450 mg	Solution: 15 mg/mL Capsule: 150 mg, 300 mg
Levofloxacin	6 months to < 5 years: 20 mg/kg/DAY divided BID ≥ 5 years: 10 mg/kg/DAY once daily	750 mg	Solution: 25 mg/mL Tablet: 250 mg, 500 mg

When to contact Pediatric Infectious Diseases (ID):

- Concern for mastoiditis or bony involvement that would require a longer duration of antibiotic therapy and monitoring
- Patients with history of and/or new cultures growing non-tuberculous *Mycobacterium* (NTM), *Actinomyces spp.*, or multi-drug resistant organisms
- Immunocompromised patients
- Complex cases in which assistance with antibiotic selection, dosing, or long-term monitoring is desired

The following is general guidance for preoperative and postoperative antibiotics in the absence of an active infection. If there is mucopurulent discharge or concern for active infection intraoperatively, please obtain appropriate cultures (aerobic, anaerobic, ± fungal, ± AFB) to guide antimicrobial therapy and contact Pediatric ID as necessary.

Procedure	Single Preoperative Antibiotic(s) Dose (if indicated)*	Postoperative Antibiotic and Duration	Ref
OTIC/EAR			
Auricular hematoma drainage	NONE	Levofloxacin x 7d while splints are in place	(1)
Ear cartilage graft	Cefazolin	Up to 24 hours (2 doses) IV cefazolin post-op	(2)
Implants (e.g., cochlear, OSIA)	Cefazolin	Up to 24 hours (2 doses) of cephalexin <i>If evidence of effusion or history of multiple ear infections within last year:</i> amoxicillin-clavulanate (clindamycin if penicillin allergy) x 5d	(2-8)
Microtia repair	Cefazolin	Continue cefazolin until drain removed, then cephalexin or amoxicillin-clavulanate (clindamycin if penicillin allergy) x 5d post-drain removal	(9)
Myringotomy	NONE	Ofloxacin ear drops x 5d	(10)
Myringoplasty	Cefazolin	NONE	(11-12)
Otoplasty	Cefazolin	NONE	(8)
Preauricular sinus surgery and/or cyst removal	Cefazolin	NONE <i>If spillage of cyst contents OR infected sinus tract</i> – Refer to Guidelines for Common Pediatric Infections: Skin and Soft Tissue Infections: Cellulitis . Antibiotic choice and duration may need adjustment based on culture results.	(13)
Tympanoplasty - with tube placement - with mastoidectomy	Cefazolin	NONE <i>If concerned for mastoiditis:</i> Refer to Guidelines for Common Pediatric Infections: Mastoiditis . Antibiotic choice and duration may need adjustment based on culture results.	(8, 14)

Procedure	Single Preoperative Antibiotic(s) Dose (if indicated)*	Postoperative Antibiotic and Duration	Ref
Salivary gland procedures (including removal/ligation) - Parotid gland and duct surgery - Salivary gland tumor removal	Cefazolin +/- metronidazole with external excision ONLY. No preoperative dose indicated for intraoral incision.	NONE * Cefazolin may be continued < 24 hours (2 doses) postoperatively for patients undergoing extensive parotid gland surgery with neck dissection	(2, 8, 15-16, 39)
Other clean otologic cases, e.g., - lesion excisions - ossicular reconstruction	NONE ‡	NONE ‡ ‡ <i>In the case of placement of new implants or with concern for low grade infection (e.g., cholesteatoma), we recommend a single dose of pre-operative cefazolin (clindamycin if penicillin allergy) be given, and post-operative antibiotics of amoxicillin-clavulanate (clindamycin if penicillin allergy) x 5d at the discretion of the surgeon.</i>	(2, 8)
Other clean-contaminated otologic cases (e.g., ear canal reconstruction)	Cefazolin		(2,17, 35)
FACIAL			
Facial fractures	NONE	NONE	(18)
Extensive facial fractures that include sinus disruption	Cefazolin	NONE If significant disruption of sinus, post-op antibiotics may be continued < 24 hours (2 doses) postoperatively.	(18)
NASAL			
Septoplasty	Cefazolin	NONE Post-operative antibiotics ONLY if splints left in place – cephalexin (clindamycin if penicillin allergy) x 7d	(2, 8, 19)
Open septorhinoplasty	Cefazolin	Autograft used: No post-operative antibiotics needed If allograft used: Post-operative antibiotics ONLY if splints placed – cephalexin (clindamycin if cephalexin allergy) x 7d	(5, 19-22)
Turbinate reduction/cautery	NONE	NONE	(8)

Procedure	Single Preoperative Antibiotic(s) Dose (if indicated)*	Postoperative Antibiotic and Duration	Ref
SINUS			
Endoscopic Sinus Surgery (e.g., antrostomy, sphenoidotomy, ethmoidectomy)	Cefazolin	NONE Post-operative antibiotics ONLY if splints left in place – cephalexin (clindamycin if penicillin allergy) x 7d	(8, 23)
OROPHARYNGEAL			
Intra-oral procedures, e.g., - adenoidectomy - mucosal/tongue biopsy - pharyngoplasty* - ranula excision - submandibular gland excision (internal approach) - tonsillectomy	NONE <i>For injection pharyngoplasty: Cefazolin</i>	NONE <i>* For injection pharyngoplasty: amoxicillin-clavulanate (clindamycin if penicillin allergy) x 7d</i>	(2, 8, 24, 39)
Dental Extraction	NONE <i>For incision through oral or pharyngeal mucosa: Cefazolin + Metronidazole</i>	<i>If extraction without concern for dental infection/abscess:</i> Third molar extractions: Amoxicillin (clindamycin if penicillin allergy) x 5d; All other: NONE <i>If extraction WITH concern for dental infection/abscess:</i> Refer to Guidelines for Common Pediatric Infections: Odontogenic infection . Antibiotic choice and duration may need to be adjusted based on culture results.	(25-27, 36-38)
Clean non-invasive scopes - Esophagoscopy - EGD - Bronchoscopy - Laryngoscopy - Sleep endoscopy (DISE)	NONE	NONE	(28-29)
Other clean-contaminated oropharyngeal cases (e.g., oral cavity resection, removal of upper airways masses)	Cefazolin + Metronidazole	NONE	(39)

Procedure	Single Preoperative Antibiotic(s) Dose (if indicated)*	Postoperative Antibiotic and Duration	Ref
CERVICAL/NECK			
Mastoidectomy	Cefazolin	NONE <i>If perioperative otorrhea or concern for infection:</i> Refer to Guidelines for Common Pediatric Infections: Mastoiditis and adjust antibiotics based on culture results	(8)
Neck mass/cyst excision and removal of sinus tract - Branchial cleft cyst - Thyroglossal duct cyst - Subglottic cyst - Submandibular gland excision (external approach)	Cefazolin	NONE <i>If spillage of cyst contents OR infected sinus tract:</i> Refer to Guidelines for Common Pediatric Infections: Cellulitis . Antibiotic choice and duration may need to be adjusted based on culture results.	(29)
Tracheoplasty	Cefazolin (no MRSA or <i>Pseudomonas</i>) Cefazolin plus vancomycin (MRSA positive) Cefepime (<i>Pseudomonas</i> positive) Cefepime plus vancomycin (MRSA and <i>Pseudomonas</i> positive)	48 hours postoperative Antibiotic based on pre-operative antibiotics	(29)
Tracheostomy/ tracheotomy	<i>Placement/Incision:</i> Cefazolin single preoperative dose <i>Tracheostomy tube removal:</i> NONE	NONE	(30-31)

Procedure	Single Preoperative Antibiotic(s) Dose (if indicated)*	Postoperative Antibiotic and Duration	Ref
Total thyroidectomy/ Parathyroidectomy	Cefazolin if surgery anticipated to last > 4hr	<i>Drain left in place:</i> cefazolin or cephalexin (clindamycin if cefazolin OR cephalexin allergy) up to 24 hours (2 doses), regardless of drain status <i>No drain:</i> NO postoperative antibiotics	(2)
Other neck surgery (clean-contaminated)	Cefazolin	NONE	(2)
CRANIOFACIAL RECONSTRUCTION			
Cranioplasty/facial reconstruction (clean)	Cefazolin	NONE	(29)
Skull base surgery Anterior, clean-contaminated Lateral, clean	Cefazolin	<i>Anterior:</i> Continue cefazolin (clindamycin if penicillin allergy) IV < 24 hours (2 doses) post-op. If > 48 hours of nasal packing/splint, use may continue cefazolin until packing is removed. <i>Lateral:</i> NONE	(2)
OTHER PLASTICS SURGERY			
Mastectomy with or without drains left in place after procedure	Cefazolin	NONE	(32-34)

*** Inpatient antibiotics per order set “ENT General Post-op”, “Microtia Reconstruction (Stage 1) Post-Op”, OR “Plastic Surgery General Post-Op”. In case of allergy, follow alternative antibiotic recommendations per order set.**

References:

1. Holtz M, Jay, Swartz J, Lu K, Donaldson R, Young N. Auricular hematoma. WikiEM. https://wikem.org/wiki/Auricular_hematoma. Updated May 8, 2017. Accessed July 24, 2024.
2. Patel PN, Jayawardena ADL, Walden RL, Penn EB, Francis DO. Evidence-Based Use of Perioperative Antibiotics in Otolaryngology. *Otolaryngol Head Neck Surg*. 2018;158(5):783-800. doi:10.1177/0194599817753610
3. Anne S, Ishman SL, Schwartz S. A Systematic Review of Perioperative Versus Prophylactic Antibiotics for Cochlear Implantation. *Ann Otol Rhinol Laryngol*. 2016;125(11):893-899. doi:10.1177/0003489416660113
4. Basavaraj S, Najaraj S, Shanks M, Wardrop P, Allen AA. Short-term versus long-term antibiotic prophylaxis in cochlear implant surgery. *Otol Neurotol*. 2004;25(5):720-722. doi:10.1097/00129492-200409000-00012
5. Hirsch BE, Blikas A, Whitaker M. Antibiotic prophylaxis in cochlear implant surgery. *Laryngoscope*. 2007;117(5):864-867. doi:10.1097/MLG.0b013e318033c2f9
6. Moon PK, Qian ZJ, Ahmad IN, Stankovic KM, Chang KW, Cheng AG. Infectious Complications Following Cochlear Implant: Risk Factors, Natural History, and Management Patterns. *Otolaryngol Head Neck Surg*. 2022 Feb 22;1945998221082530. doi: 10.1177/01945998221082530. Epub ahead of print. PMID: 35192408.
7. Kajal S, Mishra A, Gupta P, Kairo AK. Duration of Antibiotic Prophylaxis for Cochlear Implantation: A Systematic Review. *J Int Adv Otol*. 2022;18(3):269-275. doi:10.5152/iao.2022.21454
8. Ottoline AC, Tomita S, Marques Mda P, Felix F, Ferraiolo PN, Laurindo RS. Antibiotic prophylaxis in otolaryngologic surgery. *Int Arch Otorhinolaryngol*. 2013;17(1):85-91. doi:10.7162/S1809-97772013000100015
9. Eley KA, Gault DT. The bacteriology of children prior to 1st stage autologous ear reconstruction. *J Plast Reconstr Aesthet Surg*. 2010;63(12):2001-2003. doi:10.1016/j.bjps.2010.01.009
10. Alvi SA, Jones JW, Porter P, et al. Steroid Versus Antibiotic Drops in the Prevention of Postoperative Myringotomy Tube Complications. *Ann Otol Rhinol Laryngol*. 2018;127(7):445-449. doi:10.1177/0003489418776669
11. Khanal P, Guragain RP, Bhusal CL. Comparison of postoperative infection and graft uptake rate using single dose of intravenous co-amoxiclav versus no antibiotic in children undergoing myringoplasty: A randomized controlled trial. *Int J Pediatr Otorhinolaryngol*. 2020;131:109893. doi:10.1016/j.ijporl.2020.109893
12. Westman E, Höglund M, Nilsson FB, Bonnard Å, Englund E, Eriksson PO. Prophylactic antibiotics has no benefit for outcome in clean myringoplasty-A register-based cohort study from SwedEar. *Clin Otolaryngol*. 2023;48(6):895-901. doi:10.1111/coa.14089
13. Park H, Seong J, Park H, Yeo H. Standardized surgical strategy for the treatment of preauricular sinus to reduce recurrence. *Arch Craniofac Surg*. 2023;24(5):223-229. doi:10.7181/acfs.2023.00423
14. Bidkar VG, Jalisatigi RR, Naik AS, et al. Perioperative only versus extended antimicrobial usage in tympanomastoid surgery: a randomized trial. *Laryngoscope*. 2014;124(6):1459-1463. doi:10.1002/lary.24544
15. Shkedy Y, Alkan U, Roman BR, et al. Role of perioperative antibiotic treatment in parotid gland surgery. *Head Neck*. 2016;38 Suppl 1(Suppl 1):E1876-E1880. doi:10.1002/hed.24339
16. Chiesa-Estomba CM, Ninchritz E, González-García JA, Larruscain-Sarasola E, Sistiaga-Suarez JA, Altuna-Mariezcurrera X. Antibiotic Prophylaxis in Clean Head and Neck Surgery: An Observational Retrospective Single-Centre Study. *Ear Nose Throat J*. 2019;98(6):362-365. doi:10.1177/0145561319853520
17. Verschuur HP, de Wever WW, van Benthem PP. Antibiotic prophylaxis in clean and clean-contaminated ear surgery. *Cochrane Database Syst Rev*. 2004;2004(3):CD003996. doi:10.1002/14651858.CD003996.pub2
18. Goormans F, Coropciuc R, Vercruysse M, Spriet I, Willaert R, Politis C. Systemic Antibiotic Prophylaxis in Maxillofacial Trauma: A Scoping Review and Critical Appraisal. *Antibiotics (Basel)*. 2022;11(4):483. Published 2022 Apr 5. doi:10.3390/antibiotics11040483
19. Lundberg M, Lilja M, Blomgren K, et al. One dose of preoperative, intravenous, prophylactic antibiotics significantly lowers postoperative infection rate in septoplasty-a study of 772 operations. *Clin Otolaryngol*. 2022;47(1):174-180. doi:10.1111/coa.13889
20. Nandakumar G, Harb A, D'Souza AR. Grafts in septorhinoplasty: a systematic review and future directions. *Aust J Otolaryngol*. 2024;7:18.
21. Ishii LE, Tollefson TT, Basura GJ, et al. Clinical Practice Guideline: Improving Nasal Form and Function after Rhinoplasty. *Otolaryngol Head Neck Surg*. 2017;156(2_suppl):S1-S30. doi:10.1177/0194599816683153
22. Kullar, Ravina PharmD, MPH, FIDSA⁺; Frisenda, Julia MD⁺; Nassif, Paul S. MD, FACS⁺ The More the Merrier? Should Antibiotics be Used for Rhinoplasty and Septorhinoplasty?—A Review, Plastic and Reconstructive Surgery - Global Open: October 2018 - Volume 6 - Issue 10 - p e1972 doi: 10.1097/GOX.0000000000001972
23. Lehmann AE, Raquib AR, Siddiqi SH, et al. Prophylactic antibiotics after endoscopic sinus surgery for chronic rhinosinusitis: a randomized, double-blind, placebo-controlled noninferiority clinical trial. *Int Forum Allergy Rhinol*. 2021;11(7):1047-1055. doi:10.1002/alr.22756

24. Cofer SA, Baas B, Strand E, Cockerill CC. Augmentation pharyngoplasty for treatment of velopharyngeal insufficiency in children: Results with injectable dextranomer and hyaluronic acid copolymer. *Laryngoscope*. 2016;126 Suppl 8:S5-S13. doi:10.1002/lary.26227
25. Salgado-Peralvo AO, Garcia-Sanchez A, Kewalramani N, et al. Consensus Report on Preventive Antibiotic Therapy in **Dental Implant** Procedures: Summary of Recommendations from the Spanish Society of Implants. *Antibiotics (Basel)*. 2022;11(5):655. Published 2022 May 13. doi:10.3390/antibiotics11050655
26. Camps-Font O, Sábado-Bundó H, Toledano-Serrabona J, Valmaseda-de-la-Rosa N, Figueiredo R, Valmaseda-Castellón E. Antibiotic prophylaxis in the prevention of dry socket and surgical site infection after lower third molar extraction: a network meta-analysis. *Int J Oral Maxillofac Surg*. 2024;53(1):57-67. doi:10.1016/j.ijom.2023.08.001
27. Sanders JL, Houck RC. Dental **Abscess**. [Updated 2023 Feb 20]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK493149/>
28. Yetzke M, Heyes R, Nakra N, et al. Are Perioperative Antibiotics Necessary During Direct Microlaryngoscopy?. *Laryngoscope*. 2018;128(12):2838-2843. doi:10.1002/lary.27308
29. Bratzler DW, Dellinger EP, Olsen KM, et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Surg Infect (Larchmt)*. 2013;14(1):73-156. doi:10.1089/sur.2013.9999
30. Sittitrai P, Siri Wittayakorn C. Perioperative antibiotic prophylaxis in open tracheostomy: A preliminary randomized controlled trial. *Int J Surg*. 2018;54(Pt A):170-175. doi:10.1016/j.ijso.2018.04.047
31. Vander Poorten V, Uyttebroek S, Robbins KT, et al. Perioperative Antibiotics in Clean-Contaminated Head and Neck Surgery: A Systematic Review and Meta-Analysis. *Adv Ther*. 2020;37(4):1360-1380. doi:10.1007/s12325-020-01269-2
32. Bağhaki S, Soybir GR, Soran A. Guideline for Antimicrobial Prophylaxis in Breast Surgery. *J Breast Health*. 2014;10(2):79-82. Published 2014 Apr 1. doi:10.5152/tjbh.2014.1959
33. Warren DK, Peacock KM, Nickel KB, et al. Post-Discharge Prophylactic Antibiotic Following Mastectomy With and Without Breast Reconstruction. *Infect Control Hosp Epidemiol*. 2022;43(10):1382-1388. Doi:10.1017/ice.2021.400
34. Sattar AK, Masroor T, Martins RS, et al. Impact of Postoperative Antibiotic Prophylaxis on Surgical Site Infections Rates After Mastectomy with Drains but Without Immediate Reconstruction: A Multicenter, Double-Blinded, Randomized Control Superiority Trial. *Ann Surg Oncol*. 2023;30(10):5965-5973. doi:10.1245/s10434-023-13887-5
35. Vila PM, Zenga J, Jackson RS. Antibiotic Prophylaxis in Clean-Contaminated Head and Neck Surgery: A Systematic Review and Meta-analysis [published correction appears in Otolaryngol Head Neck Surg. 2018 Aug;159(2):402]. *Otolaryngol Head Neck Surg*. 2017;157(4):580-588. doi:10.1177/0194599817712215
36. Lodi G, Azzi L, Varoni EM, et al. Antibiotics to prevent complications following tooth extractions. *Cochrane Database Syst Rev*. 2021;2(2):CD003811. Published 2021 Feb 24. doi:10.1002/14651858.CD003811.pub3
37. Torof E, Morrissey H, Ball PA. The Role of Antibiotic Use in Third Molar Tooth Extractions: A Systematic Review and Meta-Analysis. *Medicina (Kaunas)*. 2023;59(3):422. Published 2023 Feb 21. doi:10.3390/medicina59030422
38. Sidana S, Mistry Y, Gandevivala A, Motwani N. Evaluation of the Need for Antibiotic Prophylaxis During Routine Intra-alveolar Dental Extractions in Healthy Patients: A Randomized Double-Blind Controlled Trial. *J Evid Based Dent Pract*. 2017;17(3):184-189. doi:10.1016/j.jebdp.2017.04.007
39. Rigotti E, Bianchini S, Nicoletti L, et al. Antimicrobial Prophylaxis in Neonates and Children Undergoing Dental, Maxillo-Facial or Ear-Nose-Throat (ENT) Surgery: A RAND/UCLA Appropriateness Method Consensus Study. *Antibiotics (Basel)*. 2022;11(3):382. Published 2022 Mar 13. doi:10.3390/antibiotics11030382